

PCT



21 DEC 1998

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: <b>A01H 4/00</b>		(11) International Publication Number: <b>WO 98/36635</b>
<b>A3</b>		(43) International Publication Date: 27 August 1998 (27.08.98)
(21) International Application Number: <b>PCT/EP98/01117</b>		(74) Agent: <b>VAN SOMEREN, Petronella, Francisca, Hendrika, Maria; Arnold &amp; Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).</b>
(22) International Filing Date: 19 February 1998 (19.02.98)		
(30) Priority Data: 97200513.6 21 February 1997 (21.02.97) EP (34) Countries for which the regional or international application was filed: NL et al. 97200701.7 10 March 1997 (10.03.97) EP (34) Countries for which the regional or international application was filed: NL et al.		
(71) Applicant (for all designated States except US): <b>VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE [BE/BE]; Rijnsschedestraat 118, Bus 1, D-9052 Zwijnaarde (BE).</b>		
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>GOETHALS, Koenraad, Hilaire, Maria [BE/BE]; Bollewerkstraat 85, B-9031 Gent (BE). EL JAZIRI, Mondher [TN/BE]; Place de la Gare 10, B-1150 Woluwe Saint Pierre (BE). VAN MONTAGU, Marc, Charles, Ernest [BE/BE]; De Stassartstraat 120, B-1050 Brussels (BE).</b>		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
		Published With international search report.
		(88) Date of publication of the international search report: 10 December 1998 (10.12.98)

## (54) Title: PLANT MICROPROPAGATION AND GERMPLASM STORAGE

Classical procedure using *in vitro* tissue culture methodUse of *Rhodococcus fascians* for micropropagating plants

## (57) Abstract

The present invention relates to a method for the *in vitro* preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of contacting plant material with a microorganism that induces fasciation and/or one or more fasciation-inducing factors derived from the microorganism or derived from the infected and/or fasciated tissue; developing leafy galls or shoot outgrowths on the plant material; and isolating the leafy galls or shoot outgrowths as the starting materials. This method may be used for the *in vitro* micropropagation of plants, which further comprises the steps of eliminating or inactivating the microorganism that induces fasciation and/or the one or more fasciation-related factors derived from the microorganism or derived from the infected and/or fasciated tissue; culturing the leafy gall or shoot outgrowths in or on one or more suitable culture media to allow shoot and root development for obtaining plantlets; and transferring the plantlets thus obtained, optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants. The method may also be used for the preservation of plants or germplasm, which is further characterized by the step of storing the leafy galls or shoot outgrowths under growth limiting conditions.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/01117

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01H4/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VEREECKE, D. ET AL: "Patterns of phenolic compounds in leafy galls of tobacco" PLANTA, vol. 201, no. 3, 1997, pages 342-348, XP002054249	1
A	CRESPI ET AL.: "Fasciation induction by phytopathogen Rhodococcus fascians depends upon a linear plasmid encoding a cytokinin synthase gene" THE EMBRO JOURNAL, vol. 11, no. 3, 1992, pages 795-804, XP002054250 cited in the application	1

---  
-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

3 September 1998

Date of mailing of the international search report

10/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tr. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fonts Cavestany, A

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/01117

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BALÁZS ET AL.: "Altered levels of Indoleacetic Acid and Cytokinin in Geranium Stems infected with C. fascians" ACTA PHYTOPATHOLOGICA ACADEMIAE SCIENTIARUM HUNGARICAE, vol. 9, no. 3-4, 1974, pages 287-292, XP002054251</p> <p>-----</p>	1

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01H</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/36635</b>
			<b>(43) International Publication Date:</b> 27 August 1998 (27.08.98)
<b>(21) International Application Number:</b> PCT/EP98/01117			<b>(74) Agent:</b> VAN SOMEREN, Petronella, Francisca, Hendrika, Maria; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).
<b>(22) International Filing Date:</b> 19 February 1998 (19.02.98)			
<b>(30) Priority Data:</b> 97200513.6 21 February 1997 (21.02.97) EP (34) Countries for which the regional or international application was filed: NL et al. 97200701.7 10 March 1997 (10.03.97) EP (34) Countries for which the regional or international application was filed: NL et al.			
<b>(71) Applicant (for all designated States except US):</b> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE [BE/BE]; Rijvisschestraat 118, Bus 1, D-9052 Zwijnaarde (BE).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GOETHALS, Koenraad, Hilaire, Maria [BE/BE]; Bollewerkstraat 85, B-9031 Gent (BE); EL JAZIRI, Mondher [TN/BE]; Place de la Gare 10, B-1150 Woluwe Saint Pierre (BE); VAN MONTAGU, Marc, Charles, Ernest [BE/BE]; De Stassartstraat 120, B-1050 Brussels (BE).			
<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).			

**Published**

Without international search report and to be republished upon receipt of that report.

**(54) Title:** PLANT MICROPROPAGATION AND GERMPLASM STORAGE**(57) Abstract**

The present invention relates to a method for the *in vitro* preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of contacting plant material with a microorganism that induces fasciation and/or one or more fasciation-inducing factors derived from the microorganism or derived from the infected and/or fasciated tissue; developing leafy galls or shoot outgrows on the plant material; and isolating the leafy galls or shoot outgrows as the starting materials. This method may be used for the *in vitro* micropropagation of plants, which further comprises the steps of eliminating or inactivating the microorganism that induces fasciation and/or the one or more fasciation-related factors derived from the microorganism or derived from the infected and/or fasciated tissue; culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow shoot and root development for obtaining plantlets; and transferring the plantlets thus obtained, optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants. The method may also be used for the preservation of plants or germplasm, which is further characterized by the step of storing the leafy galls or shoot outgrows under growth limiting conditions.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## PLANT MICROPROPAGATION AND GERMPLASM STORAGE

The present invention relates to new methods of plant micropropagation and germplasm storage.

Plants can be propagated through their two developmental life cycles: the sexual or the asexual (or vegetative). In the vegetative (asexual) cycle the unique characteristics of any individual plant selected for propagation are usually perpetuated. In most cases, each new plant produced by this method may be considered to be an extension of the somatic cell line of one individual. Methods available for propagating plants in vitro (micropropagation) are largely an extension of those already developed for conventional propagation (macropropagation).

The methods that are theoretically available for propagating plants in vitro are essentially based on the multiplication of shoots from axillary buds, on the formation of adventitious shoots and on the formation of somatic embryos. These processes are initiated either directly from the original explant or indirectly from unorganised cells such as in callus or cell suspension cultures.

In practice most micropropagated plants are produced at present by the multiplication of shoots from axillary buds. Nevertheless, somatic embryos, shoots and/or plantlets do not always originate in a single way. For example, in shoot cultures, besides axillary shoots, there are sometimes adventitious shoots formed directly on existing leave or stem explants and/or shoots arising indirectly from callus at the base of the explant.

Considering these observations, somaclonal variation is frequently observed among the regenerated plants even through multiplication of shoots from axillary buds which is considered as the most stable source for plant micropropagation. In such mixed cultures, shoots arising indirectly from callus are not desirable because of the somaclonal variation. This is particularly important when the method is applied in a large-scale programme (industrial scale). In order to

reduce this genetic/ epigenetic variation, standardization of the cellular origin for the regenerated new shoots is therefore essential.

A second limitation of the method of axillary shoot production is associated with the use of exogenous phytohormones for plant micro-propagation purposes. The growth and proliferation of axillary shoots in shoot cultures is usually promoted by incorporating growth regulators into the culture medium. Depending on the plant species, usually cytokinins and auxins in appropriate combinations and concentrations are used. Such treatments effectively remove the dominance of apical meristems so that axillary shoots are produced. These shoots are used as miniature cuttings for plant multiplication.

The cytokinin growth regulators added to shoot culture media to promote axillary shoot growth, usually inhibit root formation. Therefore, single shoots must be moved to a different medium for rooting in vitro (generally medium containing auxins) before being transferred as juvenile plantlets to the external environment. The axillary shoot production thus requires a sequence of steps.

Nowadays, there is an increased tendency to consider that exogenous applied synthetic cytokinin, which is generally too high, may cause many abnormal features (side effects) such as production of small shoots which typically fail to elongate, induction of unusual leaf shapes and particularly the tendency of the regenerated shoots to become vitrified.

In addition, high concentrations of usual cytokinins generally inhibit or considerably delay root formation and also prevent root growth and the promoting effects of auxins on root initiation. In this respect, frequently more than one subculture to a cytokinin-free medium may be required until the level of cytokinin within the tissues has been sufficiently reduced followed by a transfer of the shoots to a rooting medium.

Considering these aspects it is furthermore impossible to propagate particular plant species which are sensitive to such hormonal treatments.

It is thus the object of the present invention to  
5 provide an alternative micropropagation method with which the above identified drawbacks of the known methods are obviated.

According to the invention it has surprisingly been found that leafy galls, which are produced on  
10 dicotyledonous plants by infection with particular micro-organisms or by treatment with substances produced by these micro-organisms or derived from the infected and/or fasciated tissue (so-called fasciation), and which consist of multiple meristematic centers and shoot  
15 primordia that are suppressed for elongation, are an excellent starting point for micropropagation. In a gall no apical dominance phenomena are evident and the formation of numerous primordia is derepressed by the infection process. The gall is in fact a large mass of  
20 shoot primordia that are covered with leaves and that are infected by the micro-organism. In leafy galls specific metabolites are present that are involved in the establishment and the maintenance of the leafy gall structure.

25 In monocotyledonous plants the structures that arise upon contact with or infection by fasciation-inducing micro-organisms or substances have a different morphology than the above described leafy galls that are formed in dicotyledonous plant material. Infection either  
30 results in deformation of bulbs and/or the formation of several elongated shoots. These monocot elongated shoot structures are useful in the method of the invention and will be further identified as "shoot outgrows", whereas the dicot structures are called "leafy galls".

35 Apart from the morphogenetic aspects, a gall represents a center of elongation inhibition where the bacterium suppresses the outgrowth of the primordia in the gall. This phenomenon is translated in the apparent



lack of apical dominance that would be exerted by one of the primordia and would lead to the outgrowth of distinct shoot primordia from the gall tissues. As such, each gall can be considered as a highly condensed plant where the amplified production of primordia and meristems -apical as well as axillary- are confined to one compact region through the suppression of both the elongation processes and apical dominance.

Furthermore, it was found according to the invention that due to the fact that the galls consist of large numbers of shoot primordia confined to a compact and small structure, they are also a very suitable material for germplasm storage.

Starting from these two observations the present invention now provides for a method for the in vitro preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of:

- a) contacting plant material with a micro-organism that induces fasciation and/or one or more fasciation-inducing factors derived from the micro-organism, or derived from the infected and/or fasciated tissue;
- b) developing leafy galls or shoot outgrows on the plant material; and
- c) isolating the leafy galls or shoot outgrows as the starting materials.

This method of the invention may then be incorporated in a method of micropropagation or a method of germplasm storage, respectively. A method for the in vitro micropropagation of plants comprises the above described method for the preparation of starting material, further characterized by the steps of

- d) eliminating or inactivating the micro-organism that induces fasciation and/or the one or more fasciation-related factors derived from the micro-organism or derived from the infected and/or fasciated tissue;

e) culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow further shoot and root development for obtaining plantlets, and

- 5 f) transferring the plantlets thus obtained, optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants.

A method for the preservation of plants or germplasm comprises the method for the preparation of  
10 starting material, further characterized by the step of storing the leafy galls or shoot outgrows under growth limiting conditions. The growth limiting conditions may comprise low temperatures of about 4°C or cryo-conserving conditions, such as storage in liquid nitrogen.

- 15 Both methods are not only applicable on angiosperms, but also on gymnosperms.

In a preferred embodiment of the invention the fasciation inducing micro-organism is the Gram positive bacterium Rhodococcus fascians (Tilford) (Goodfellow M. &  
20 Cross, T., "Classification", published: Goodfellow, Modarski & Williams ed., The Biology of Actinomycetes, Academic Press London, p. 7-64 (1984)). (Effective publication Goodfellow M., "Reclassification of Corynebacterium fascians (Tilford, Dowson) in the genus  
25 Rhodococcus, as Rhodococcus fasc. comb. nov." System. Appl. Microbiol. 5, 225-229 (1984), previously known as Phytomonas fascians Tilford, J., Agric. Res. 53, 383-394 (1936); or Corynebacterium fascians (Tilford) Dowson, W.J., Trans. Brit. Myc. Soc. 25, 311-314 (1942)).  
30 Rhodococcus fascians is a Nocardioform Actinomycete that infects a whole range of both dicotyledonous and monocotyledonous plants.

The range of plants that can be infected by R. fascians is very broad with different species belonging  
35 to numerous plant families, such as Brassicaceae, Salicaceae, Compositae, Solanaceae, Scrophulariaceae, Liliaceae, Papaveraceae, Gramineae and Fabaceae.

Infection of an intact plant can for example be achieved by vacuum infiltration of plants with suspensions of the bacteria, application of a dense bacterial culture sample to a wound surface or, less efficiently, by simply applying bacteria in either the growth substrate of the plants or on the surface of intact plants. Also explants, such as leaf discs, organs, tissue fragments, and seedlings, seeds, embryos, isolated cells and protoplasts can be successfully infected, giving rise to leafy galls or shoot outgrows. Furthermore, R. fascians infected cell cultures and callus tissues react in a similar fashion.

The role of R. fascians in the phenomenon of suppression of elongation and apical dominance can be exemplified by killing of the bacteria in the gall through a treatment with bactericidal agents such as Carbenicillin, Triacellin and Claforan. Growth of detached galls on hormone free plant media with the addition of bactericidal compounds to an end concentration of 500 mg/L, results in the release of the amplified primordia from the bacterial suppression. ensuing in the outgrowth of numerous independent shoots from the gall tissues that will produce root tissues. The rooting can be facilitated by treating the galls in the dark where the release of the shoots is accompanied by etiolation, producing masses of elongated sprouts. Each shoot can be transferred to another medium resulting in the production of a fertile plant clone.

These new micropropagation and germplasm storage methods are widely applicable and may for example be used for trees, crops, medicinal and aromatic plants. The methods are also suitable for biochemical or genetic selection purposes.

Instead of using the micro-organism as a whole one or more factors derived from the micro-organism or infected and/or fasciated tissue may be used. These factors are capable of inducing the formation of leafy galls in dicotyledonous plants and shoot outgrows in

monocotyledonous plants. The factors may for example be gene products of the genes involved in fasciation, or the molecules produced by these gene products.

In the R. fascians strain D188, the property  
5 to induce fasciation is correlated with the presence of a  
+200 kb large linear plasmid, pFiD188, that carries  
several loci (e.g. fas, att and hyp) involved in leafy  
gall formation. The present inventors have found that  
mutations in the fas locus render the bacteria non-  
10 pathogenic. A mutation in the att locus provokes  
attenuated fasciation resulting in smaller leafy galls  
that are formed more slowly. Finally, a mutation in the  
hyp locus results in hyper-fasciation, leading to the  
formation of larger galls compared to wild type.

15 Apart from pFiD188 encoded genes, also chromosomal  
genes in the bacterium are involved in the infection  
process for instance by influencing the endophytic/  
epiphytic survival of R. fascians.

Evidently, gall formation results from the  
20 modulation of the regular hormone balance in the infected  
plant tissues. Although the fasciation symptoms  
superficially resemble effects caused by the application  
of different doses of several cytokinins, they are  
nevertheless clearly discriminated from the latter by the  
25 presence of non-elongating primordia and the absence of  
abundantly developing shoots. Early reports where galling  
was correlated with the presence of different cytokinins  
in the growth medium of R. fascians have recently been  
questioned by opposing results. This is further  
30 substantiated by the present inventors by their finding  
of the presence of cytokinins in the growth media of a  
non-fasciating strain D188-5 (Crespi et al., EMBO J. 11,  
795-804 (1992)) cured for pFiD188.

Although early results hinted at the presence of  
35 an adenosine monophosphate isopentenyl transferase (ipt)  
homologous gene in the fas locus on pFiD188 that would be  
involved in the biosynthesis of cytokinins (Crespi et  
al., 1992, supra), the present results question the

unique involvement of the ipt homologue in the biosynthesis of regular cytokinins like isopentenyl adenine or zeatin. Instead, the fas locus is involved in the production and secretion of one or more

5 isopentenylated molecules that affect the plants hormone levels. This was confirmed by the isolation and preliminary chemical analysis of one fas dependent products and fits with the overall low amino acid identity between the fas encoded ipt gene product and

10 regular isopentenyl transferase proteins. Apart from the ipt related gene, the fas locus encodes several genes that, on the basis of sequence homology studies, are implicated in cytochrome P450 dependent oxygenation and S-Adenosyl Methionine dependent methylation processes

15 of, conceivably, the fas product.

High expression of the fas locus in the bacterium -as measured by translational GUS fusion assays- is strictly controlled and can only be consistently achieved in batch cultures using a defused

20 induction medium, such as Minimal A-medium (obtainable by adding 20.5 ml 0.1M citrate and 29.5 ml 0.1M Na-citrate to 450 ml distilled water and adding after autoclaving and cooling 500 $\mu$ l 1% (w/v) thiamine, 1.25 ml MgSO<sub>4</sub> and 1.25 ml 40% NH<sub>4</sub>SO<sub>4</sub>) and only in the presence of either

25 gall extracts or histidine in combination with particular carbon sources such as pyruvate and succinate. Plant extracts cannot induce fas gene expression, but gall extracts contain a specific inducer molecule. This inducer molecule is present in all gall tissues formed on a whole

30 range of different plants and is able to induce expression in minimal concentrations. Furthermore, production of this molecule was shown in tobacco BY-2 cell cultures confronted with R. fascians wild type cells while this was not the case for non-infected plant cell

35 cultures or pure bacterial cultures.

The fas expression is mainly regulated at the post-transcriptional level, indirectly through the activity of the fasR gene that encodes an AraC type

transcriptional activator and possibly directly through the hyp locus that carries a gene encoding an RNA-helicase.

Confrontation of BY-2 tobacco cell culture  
5 synchronized for mitosis using aphidicolin with fas  
induced D188 cells resulted in a marked increase of the number of tobacco cells that enter the first round of mitosis upon relieve of the mitosis block. This effect is absent when the tobacco cell culture is confronted with  
10 the non-fasciating strain D188-5 or with fas mutant strains.

Classical cytokinin-bioassays using extracts of R. fascians D188 and D188-5 grown for fas induction proved difficult to interpret due to the presence of high  
15 background levels of contaminating cytokinins in both tested extracts. From D188 cultures, induced for fas expression, a fas dependent metabolite was isolated.

The att locus comprises a 20 kb region of pFiD188, linked to the fas locus. Sequencing showed the  
20 presence of several genes that are involved in the synthesis and secretion of a complex molecule of unknown structure. Apart from genes involved in the synthesis of arginine, the locus carries genes implicated in the biosynthesis and modification of fatty acid related  
25 molecules and a gene related to export protein genes.

Expression of the att locus is partially co-regulated with the fas locus; induction of GUS fusions is only visible in defined media in the presence of gall extracts or of histidine in combination with particular  
30 carbon sources and depends on the activity of the fasR gene. However, an independent regulation through an att specific regulatory gene, attR, is postulated here whereby the expression of the attR gene is itself enhanced by the fasR gene in response to plant extracts.

35 Using <sup>14</sup>C-labelled arginine or acetate, in either case the induced accumulation in the growth medium of R. fascians D188 of the same <sup>14</sup>C-labelled, att dependent compounds was evident using thin layer chromatography,

proving the incorporation of both tested substrates in complex, secreted compounds. These were isolated through High Pressure Liquid Chromatography for chemical analysis and bio-assaying using BY-2 tobacco cell cultures. As an example, a molecule with a molecular weight of 681 dalton was isolated.

The products of the att locus are probably involved in the formation of leafy galls in combination with the fas locus as a mutant in the locus is strongly retarded in the galling process. It is also conceivable that the att and fas loci cooperate in the biosynthesis and secretion of a single type of molecules that provoke fasciation.

Apart from the fas and att loci, still other loci on pFiD188 are involved in the formation of leafy galls. The hyp locus probably negatively controls expression of the fas and att expression through RNA helicase activity on the mRNA. Further downstream of the fas locus another locus has been isolated that is essential for fasciation and sequencing of which showed the presence of oligopeptide synthetic genes.

Molecular analysis of the gene expression in infected tobacco plants using differential display techniques showed the induction of different genes showing homology to genes involved in the metabolism of gibberellins and steroids as well as the repression of a gene showing homology to oligopeptide receptor proteins. These results strongly suggest an influence of the infecting bacteria and their products on the existing balances of different plant growth factors, resulting in the induction of meristem and primordium formation.

The formation of a leafy gall by R. fascians is thus a complex process with different levels of signalling between both partners. The expression of the bacterial fasciation genes located on the linear plasmid is prone to a complex regulation that ultimately implies the involvement of a gall specific metabolite in the fasR dependent induction of the att and fas genes. Once the

fas and att genes are optimally expressed, metabolites are produced and secreted in the plant that disturb and/or redirect the hormone household of the plant tissues. The fas and att derived factor(s) are implicated in the enhancement of plant cell divisions and are crucial for the activation of the gall formation. Molecular data suggest the involvement of gibberellins, steroid type molecules and/or possibly oligopeptides of the plant. This unique response of the plant to infection confers the particular properties of R. fascians and its derived products in altering the developmental programming of the plant in some hitherto non-described way. It further discriminates the leafy gall syndrome from other neoplastic pathologies like the Agrobacterium tumefaciens induced crown gall formation and the Pseudomonas savastanoi induced galling of Oleander species.

In summary, it has now been found that a gall constitutes several meristems and primordia the formation of which is correlated with the fas product, the att products, or a combination of both and possibly other products the synthesis of which is determined by different loci on pFiD188 or on the bacterial chromosome. The end result is a local activation and deregulation of primordium and meristem formation that results in dense tissues -the gall- that is ultimately covered with leaflets. Morphological analysis of these galls suggest an organised spatial distribution of the primordia that are covered with leaflets, both according to the fundamental rules of phyllotaxis.

Based on the experimental information provided here it became possible to identify a complex of factors that can be used to induce fasciation independently from the bacteria as a whole. This complex of factors comprises at least the fas products and/or the att products and in addition other gene products encoded by the pFiD188 plasmid.



The unique property of R. fascians to amplify shoot primordium formation through the modulation of specific hormone balances in a broad spectrum of plants is useful for the establishment of a regeneration program for a diverse range of plants as will be illustrated in the following examples. Infection of plants with R. fascians leads to the local amplification of shoot primordia, combined with a suppression of further shoot growth and development. Upon amplification, primordia can be released from the suppressive effects of R. fascians by killing of the bacteria. This can be achieved by detaching galls and growing them on media with doses of bactericides like penicillin's. Any other treatment of the galls that will kill the inciting bacteria without harming the plant tissue can be envisioned here.

As a result, from the gall tissues shoots will sprout that can be detached and that will form roots, ultimately giving rise to normal, regenerated plants. This effect is even more dramatic when the galls are treated in the dark, where the release of shoots is accompanied by etiolation, producing masses of elongated sprouts that can be propagated to a normal plant.

This technique has been successfully applied to leafy galls on Nicotiana tabacum, Papaver somniferum, Artemisia annua, Vigna unguiculata, Atropa belladonna, Sesbania rostrata and Digitalis lanata.

The Figure is an overview showing the steps required for micropropagation using either classical procedure or Rhodococcus fascians.

This invention as described here will be further illustrated in the following examples.

## EXAMPLES

### EXAMPLE 1

Regeneration of plants from leafy galls of various plant species-induced by Rhodococcus fascians

5

#### 1. General method

The following method is applicable to plant species in general. In these examples the plant species Nicotiana tabacum, Papaver somniferum, Artemisia annua,  
10 Vigna unguiculata, Atropa belladonna, Sesbania rostrata and Digitalis lanata were used.

##### 1.1. Bacterial strain

Rhodococcus fascians strain D188 or other  
15 virulent strains are cultured in solid or liquid conventional bacteria medium or a recommended one containing: Glucose (10 g/l), Yeast extract (5 g/l), Peptone (5 g/l), Agar (15 g/l), distilled water 1l, the pH adjusted to 7.0. The cultures are then incubated in  
20 the dark at 25°C.

##### 1.2. Plant material

Selected plant species are used for the propagation purposes. After surface disinfection of the  
25 explants (for example: cutting consisting of stem internode fragments with axillary buds) using appropriate reagents known in the art, the explant is transferred to in vitro culture condition using conventional tissue culture medium such as the well-known basal Murashige and  
30 Skoog medium (Physiol. Plant. 15, 473-497) or Gamborg medium (Gamborg, O.L. et al., Ex. Cell. Res. 50, 151 (1986)).

##### 1.3. Infection procedure

35 The axenic explant is directly infected with the fast growing bacteria or infected under vacuum. All other method of infection which make close contact of the bacteria and the plant tissue or cells are also suitable.

In case of Sesbania rostrata, cotyledonous explants were used for direct infection with Rhodococcus fascians. Preferably wounded cotyledons were used for this infection in order to obtain a more efficient infection.

5

#### 1.4. Incubation of the infected explant

Containers containing the infected plant material are incubated in the room culture at 23°C. Depending on the plant species, leafy galls came to be visible after 10 3-4 weeks of incubation.

### 2. Regeneration of plants Leafy galls structure

#### 2.1. General procedure

The leafy galls are first isolated from the 15 original explants. Then they are subcultured in a hormone-free conventional plant tissue culture medium additionated with bactericide reagent such as Claforan at final concentration of 500 mg/l. The cultures are preferably incubated in the dark at appropriate 20 temperature (depending on the plant species, for example 19°C for Papaveraceae species; 23°C for Solanaceae species).

#### 2.2. Results

25 The dark condition combined with the antibiotic treatment improve the shoot elongation. The etiolated shoots thus obtained are then subcultured in a hormone-free culture medium and incubated in light condition. The light intensity as well as the temperature of 30 incubation are selected according to the specific requirement of each plant species under investigation. After several days of cultures, spontaneous root appeared, and thus plantlet are recovered.

Using the described method and in the case of 35 Digitalis lanata, an average of 60 new regenerated shoots/explant are obtained instead of 12 new regenerated shoots using the classical micropropagation method (using exogenous cytokinins).

Using common method for acclimatization, these new regenerated plantlets are successfully transferred to the green house.

Quantitative and qualitative analysis of cardenolides in the new regenerated plants showed a distribution quite similar to that of the mother plant. In addition the morphological traits of these new regenerated plants are similar to those of the mother plant, indicating a reduced variability as compared to the classical micropropagation technique.

## EXAMPLE 2

### Germplasm storage

#### 1. Short-term conservation

The leafy galls containing the bacteria are isolated from the original explant then subcultured in a conventional plant tissue culture medium with reduced sugar concentration (15g/l of sucrose). After one week of culture at 20°C, the culture containers are transferred to 4°C under dark condition. In such conditions the capability of the shoots to develop (after bactericide treatment) is maintained for 1-2 years.

#### 2. Long term conservation

The well established procedure for cryo-conservation using liquid nitrogen is also convenient for the leafy gall structure (George E.F., in: Plant Propagation by Tissue Culture, Part I, George Edwin F. (Ed.), Exegetics Ltd, Edington, Wilts, England, pp. 163-181 (1993)).

## CLAIMS

1. Method for the in vitro preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of:

a) contacting plant material with a micro-  
5 organism that induces fasciation and/or one or more fasciation-inducing factors derived from the micro-organism or derived from the infected and/or fasciated tissue;

b) developing leafy galls or shoot outgrows on  
10 the plant material; and

c) isolating the leafy galls or shoot outgrows as the starting materials.

2. Method for the in vitro micropropagation of plants, which method comprises the method as claimed in  
15 claim 1, further characterized by the steps of

d) eliminating or inactivating the micro-organism that induces fasciation and/or the one or more fasciation-related factors derived from the micro-organism or derived from the infected and/or fasciated  
20 tissue;

e) culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow shoot and root development for obtaining plantlets; and

f) transferring the plantlets thus obtained,  
25 optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants.

3. Method for the preservation of plants or germplasm, which method comprises the method as claimed in claim 1, further characterized by the step of storing  
30 the leafy galls or shoot outgrows under growth limiting conditions.

4. Method as claimed in claims 1-3, wherein the fasciation-inducing micro-organism is Rhodococcus fascians.

35 5. Method as claimed in claim 3 and 4, wherein the growth limiting conditions comprise low temperatures


of about 4°C or cryo-conserving conditions, such as storage in liquid nitrogen.

6. Method as claimed in claims 1-5, wherein the plant material comprises explants, such as leaf discs, 5 organs, tissue fragments, seedlings, seeds, embryos, isolated cells, protoplasts, cell cultures and callus tissues.

7. Method as claimed in claims 1-6, wherein the plant material is material of species of the genera 10 Brassicaceae, Salicaceae, Compositae, Solanaceae, Scrophulariaceae, Liliaceae, Papaveraceae, Gramineae and Fabaceae.

8. Method for the in vitro micropropagation of plants, which method comprises the method of claim 3 and 15 optionally the subsequent removal or inactivation of still present micro-organisms that induce fasciation and/or fasciation related factors derived from the micro-organism or derived from the infected and/or fasciated tissue, culturing the leafy gall or shoot outgrows in or 20 on one or more suitable culture media to induce shoot and root development for obtaining plantlets, and transferring the plantlets thus obtained, optionally after acclimisation, to conventional growing conditions to obtain regenerated plants.

Classical procedure using in vitro tissue culture method



Use of *Rhodococcus fascians* for micropropagating plants

